

J. Balasubramaniam[†] and G V V Sharma^{*}

Abstract—This manual covers the properties of sequences through examples. Python scripts are provided for understanding the properties of sequences.

1 LIMIT

Problem 1.1. Sketch the following sequence.

$$x_n = \frac{2n}{n + 4\sqrt{n}}, \quad n \geq 0 \quad (1.1)$$

Solution:

```
import numpy as np
import matplotlib.pyplot as plt

n = np.linspace(0,1e9,100)
T_n = (2.0*n)/(n+4.0*np.sqrt(n))
plt.plot(n,T_n)
plt.grid()
plt.xlabel('$n$')
plt.ylabel('$x_n$')
#Comment the following line
#plt.savefig('../figs/seq_converge
    .eps')
plt.show()
```

Definition 1.1. The sequence x_n converges to a limit L if for $\epsilon > 0$, there exists an integer $K(\epsilon)$ such that for all $n > K(\epsilon)$,

$$|x_n - L| < \epsilon \quad (1.2)$$

Proposition 1.1. Archimedian Property: For any real number x , there exists an integer $n > x$.

Problem 1.2. Guess the value of L for x_n in (1.1) as $n \rightarrow \infty$.

[†] The author is with the Department of Mathematics, IIT Hyderabad. ^{*}The author is with the Department of Electrical Engineering, IIT, Hyderabad 502285 India e-mail: {jbala,gadepall}@iith.ac.in. All material in the manuscript is released under GNU GPL. Free to use for all.




Fig. 1.1

Solution: From Fig. 1.1, $L = 2$.

Problem 1.3. Show that x_n in (1.1) converges to $L = 2$ using Definition 1.1.

Solution:

$$|x_n - 2| = \left| \frac{2n}{n + 4\sqrt{n}} - 2 \right| \quad (1.3)$$

$$= \frac{8\sqrt{n}}{n + 4\sqrt{n}} \quad (1.4)$$

$$< \frac{8\sqrt{n}}{n} = \frac{8}{\sqrt{n}} \quad (1.5)$$

Using Proposition 1.1, choose $K(\epsilon) > \frac{64}{\epsilon^2}$ to be an integer.

$$n > K(\epsilon) \Rightarrow n > \frac{64}{\epsilon^2} \Rightarrow \frac{8}{\sqrt{n}} < \epsilon \quad (1.6)$$

Thus, there exists $K(\epsilon)$ such that $|x_n - 2| < \epsilon$.

Problem 1.4. Let $x_n = \frac{1}{\ln(n+1)}$.

- 1) Find the value to which x_n converges.
- 2) Find $K(\epsilon)$ when $\epsilon = \frac{1}{2}$ and $\epsilon = \frac{1}{10}$.

2 MONOTONICITY AND BOUNDEDNESS

Definition 2.1. A sequence x_n is said to be monotonically *increasing* if

$$x_{n+1} > x_n \quad (2.1)$$

x_n is monotonically *decreasing* if $x_{n+1} < x_n$.

Definition 2.2. The sequence x_n is said to be bounded if for all $n > N$

$$|x_n| < M \quad (2.2)$$

for some positive real number M .

Problem 2.1. Consider the sequence defined by

$$x_n = \begin{cases} 1 & n = 1 \\ \frac{x_{n-1} + 1}{3} & n > 1 \end{cases} \quad (2.3)$$

Is the sequence

- 1) Monotonic?
- 2) Bounded?

Solution: The following code plots Fig. 2.1. It is obvious from the figure that x_n is both monotonically decreasing as well as bounded.

```
import numpy as np
import matplotlib.pyplot as plt

x = []
temp = 1
for i in range(100):
    x.append(temp)
    temp = (temp + 1.0) / 3.0

plt.plot(range(100), x)
plt.grid()
plt.xlabel('$n$')
plt.ylabel('$x_n$')
#Comment the following line
#plt.savefig('../figs/seq_monotone
    .eps')
plt.show()
```

Problem 2.2. Prove that x_n is monotonically decreasing.

Proof. From (2.3), it can be shown that

$$x_n = \frac{1}{2} \left(1 + \frac{1}{3^{n-1}} \right) \quad (2.4)$$




Fig. 2.1

Thus,

$$x_{n-1} - x_n = \frac{2}{3^{n-1}} > 0 \Rightarrow x_{n-1} > x_n \quad (2.5)$$

which is the condition for x_n to be monotonically decreasing.

Problem 2.3. Prove that x_n is monotonically decreasing using induction.

Proof. Since $x_1 = 1, x_2 = \frac{2}{3} < x_1$. Let $x_k < x_{k-1}$. Then

$$x_{k+1} - x_k = \frac{x_k + 1}{3} - \frac{x_{k-1} + 1}{3} \quad (2.6)$$

$$= \frac{x_k - x_{k-1}}{3} < 0 \quad (2.7)$$

Thus, $x_k < x_{k-1} \Rightarrow x_{k+1} < x_k$. This shows that x_n is decreasing.

Problem 2.4. Show that x_n is bounded.

Proof. From (2.4), it is obvious that

$$|x_n| \leq 1 \quad (2.8)$$

Thus, x_n is bounded. \square

Problem 2.5. Find the limit of x_n .

Solution: From Fig. 2.1, it is clear that the limit is $\frac{1}{2}$.

Problem 2.6. Show that the limit of x_n is $\frac{1}{2}$.

Problem 2.7. Show that the sequence defined by

$$x_n = \begin{cases} 2 & n = 1 \\ \sqrt{2x_{n-1} + 1} & n > 1 \end{cases} \quad (2.9)$$

is monotone as well as bounded. Find its limit.

Proposition 2.1. Any monotone sequence that is bounded is convergent.

Problem 2.8. Graphically show that $x_n = \sqrt{n+1}-1$ is divergent.

Solution: The following code results in Fig. 2.8. It is obvious that the series does not converge.

```
import numpy as np
import matplotlib.pyplot as plt

n = np.linspace(0,1e3,100)
x_n = np.sqrt(n+1) - 1
plt.plot(n,x_n)
plt.grid()
plt.xlabel('$n$')
plt.ylabel('$x_n$')
#Comment the following line
#plt.savefig('../figs/seq_diverge.
eps')
plt.show()
```




Fig. 2.8

Problem 2.9. Show that x_n in Problem 2.8 is increasing.

Proof. Since $x_n > x_{n-1}$ for all n , the sequence is increasing. \square

Problem 2.10. Show that x_n in Problem 2.8 is unbounded.

Proof. For every M , it is possible to find an integer $n > M^2 - 1 \Rightarrow \sqrt{n+1} > M$. Thus, x_n is unbounded. \square

Proposition 2.2. A monotone unbounded sequence is divergent.

3 CAUCHY SEQUENCE

Definition 3.1. The sequence x_n is Cauchy if for every $\epsilon > 0$, there exists an integer N such that

$$|x_m - x_n| < \epsilon \quad \text{whenever } n, m \geq N \quad (3.1)$$

Problem 3.1. Show that

$$x_n = \frac{1}{n^2} \quad (3.2)$$

is a Cauchy sequence.

Proof. Let $m > n > N$.

$$|x_m - x_n| = \left| \frac{1}{m^2} - \frac{1}{n^2} \right| = \left| \frac{(m-n)(m+n)}{m^2 n^2} \right| \quad (3.3)$$

The numerator in (3.3)

$$|(m-n)(m+n)| < 2m^2 \quad (3.4)$$

Thus,

$$|x_m - x_n| < \frac{2}{n^2} < \frac{2}{N^2} < \epsilon \quad (3.5)$$

Thus it is possible to find an N given ϵ for x_n such that x_n is Cauchy.

Proposition 3.1. Every Cauchy sequence is convergent and vice versa.

Problem 3.2. Show that

$$x_n = 1 + \frac{1}{2!} + \frac{1}{3!} \cdots + \frac{1}{n!} \quad (3.6)$$

is a Cauchy sequence.

Problem 3.3. Is $x_n = \sqrt{n}$ a Cauchy sequence?