

G V V Sharma*

CONTENTS

| | | |
|----------|---|----------|
| 1 | Medians of a Triangle | 1 |
| 1.1 | Coordinates of Point Dividing a Line in Ratio $k : 1$. . . | 1 |
| 1.2 | Median | 2 |
| 2 | Altitudes of a Triangle | 4 |
| 2.1 | Lines and Angles | 4 |
| 2.2 | Altitude | 5 |
| 3 | The Circle | 6 |
| 3.1 | Definitions | 6 |
| 3.2 | Tangent | 6 |

Abstract—The focus of this text is on investigating the properties of triangles and circles through coordinate geometry. An early exposure to coordinate geometry allows students to use computers for mathematical visualizations and computing. This book provides an alternative approach to geometry which may be useful to those having difficulties with traditional methods. In the process, almost all the basic concepts of coordinate geometry are covered. Instead of teaching geometry and coordinate geometry as separate subjects in high school, this textbook shows how to develop a holistic approach for teaching math.

1 MEDIANS OF A TRIANGLE

1.1 Coordinates of Point Dividing a Line in Ratio $k : 1$

Definition 1.1. (Budhayana’s Theorem:)In Fig. 1.1, ABC is a right angled triangle with $AB = b$ and $BC = c$. Using coordinates, A, B and C can be represented as $(a, b), (0, 0)$ and $(c, 0)$, where B is the origin, BC is on the X -axis and AB is on the Y -axis. Using Budhayana’s theorem, $AC = \sqrt{b^2 + c^2}$.

*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in.




Fig. 1.1: Budhayana’s Theorem

Problem 1.2. Let A and B have the coordinates (a_1, a_2) and (b_1, b_2) respectively. Show that the distance between A and B is given by

$$AB = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2} \quad (1.1)$$

Proof. In Fig. 1.2.1, it is obvious that

$$PB = A_1B_1 = a_1 - b_1 \quad (1.2)$$

$$PA = A_2B_2 = a_2 - b_2 \quad (1.3)$$

Using Budhayana’s theorem,

$$AB = \sqrt{PA^2 + PB^2} = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2} \quad (1.4)$$

An alternative visualization of Fig. 1.2.1 is available in 1.2.2, where the line AB is shifted to the origin such that the line becomes $A'O$. This results in

$$OA' = \sqrt{(A'P')^2 + (O'P')^2} = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2} \quad (1.5)$$

Thus, shifting the coordinates by (b_1, b_2) makes the visualization as well as computation easier. This trick will be used quite a lot in this book.




Fig. 1.2.1: Distance formula.





Fig. 1.2.2: Alternative visualization of distance.

Problem 1.3. In Fig. 1.3, P divides AB in the ratio $k : 1$ such that

$$\frac{PA}{PB} = k. \quad (1.6)$$

If the coordinates of A and B are (a_1, a_2) and (b_1, b_2) respectively, show that the coordinates of P are

$$\left(\frac{a_1 + kb_1}{k+1}, \frac{a_2 + kb_2}{k+1} \right) \quad (1.7)$$

Fig. 1.3: P divides AB in the ratio $k : 1$.

Proof. From Fig. 1.3, using similar triangles, it is obvious that

$$\frac{PB}{PA} = \frac{P_1B_1}{P_1A_1} = \frac{x - b_1}{a_1 - x} = k \quad (1.8)$$

Simplifying the above,

$$x - b_1 = k(a_1 - x) \quad (1.9)$$

$$\Rightarrow (1 + k)x = ka_1 + b_1 \quad (1.10)$$

$$\Rightarrow x = \frac{ka_1 + b_1}{k + 1} \quad (1.11)$$

The proof for y is similar.

1.2 Median

Definition 1.4. The line segment AD in Fig. 1.4 that divides the side BC in two equal halves ($BD = DC$) is known as the median.




Fig. 1.4: Median

Problem 1.5. The medians BE and CF in Fig. 1.5 intersect at O , such that

$$\begin{aligned} \frac{OB}{OE} &= k_1 \\ \frac{OC}{OF} &= k_2 \end{aligned} \quad (1.12)$$

Show that $k_1 = k_2 = 2$.




Fig. 1.5: Medians BE and CF

Proof. Let the coordinates of A , B and C be (a, b) , $(0, 0)$ and $(c, 0)$ respectively. Using 1.7,

$$E = \left(\frac{a+c}{2}, \frac{b}{2} \right) \quad (1.13)$$

$$F = \left(\frac{a}{2}, \frac{b}{2} \right) \quad (1.14)$$

Since O divides BE in the ratio $k_1 : 1$, from (1.7),

$$O = \frac{k_1 E + B}{k_1 + 1} = \frac{1}{k_1 + 1} \left[k_1 \left(\frac{a+c}{2}, \frac{b}{2} \right) + \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right] \quad (1.15)$$

$$= \frac{k_1}{2(k_1 + 1)} \begin{pmatrix} a+c \\ b \end{pmatrix} \quad (1.16)$$

Similarly, since O divides CF in the ratio $k_2 : 1$,

$$O = \frac{k_2 F + C}{k_2 + 1} = \frac{1}{k_2 + 1} \left[k_2 \left(\frac{a}{2}, \frac{b}{2} \right) + \begin{pmatrix} c \\ 0 \end{pmatrix} \right] \quad (1.17)$$

$$= \frac{1}{2(k_2 + 1)} \begin{pmatrix} ak_2 + 2c \\ bk_2 \end{pmatrix} \quad (1.18)$$

From (1.16) and (1.18),

$$\frac{k_1}{2(k_1 + 1)} \begin{pmatrix} a+c \\ b \end{pmatrix} = \frac{1}{2(k_2 + 1)} \begin{pmatrix} ak_2 + 2c \\ bk_2 \end{pmatrix} \quad (1.19)$$

$$\Rightarrow \frac{k_1}{(k_1 + 1)} \begin{pmatrix} a+c \\ b \end{pmatrix} = \frac{1}{(k_2 + 1)} \begin{pmatrix} ak_2 + 2c \\ bk_2 \end{pmatrix} \quad (1.20)$$

By equating the respective coordinates,

$$\frac{k_1(a+c)}{k_1+1} = \frac{ak_2+2c}{k_2+1} \quad (1.21)$$

$$\frac{k_1 b}{k_1+1} = \frac{k_2 b}{k_2+1} \quad (1.22)$$

From (1.22), it is trivial to obtain $k_1 = k_2$. Substituting this in (1.21) and simplifying,

$$k_1(a+c) = ak_1 + 2c \quad (1.23)$$

$$\Rightarrow k_1 = 2 \quad (1.24)$$

Thus $k_1 = k_2 = 2$.

Problem 1.6. In Fig.1.6,

$$\frac{OB}{OE} = \frac{OC}{OF} = 2 \quad (1.25)$$

Show that BE and CF are medians.




Fig. 1.6: O divides BE and CF in the ratio $2 : 1$.

Proof. Let E divide AC in the ratio $k_2 : 1$ and F divide AB in the ratio $k_1 : 1$. Using (1.7),

$$E = \frac{k_2 C + A}{k_2 + 1} = \frac{1}{k_2 + 1} \left[k_2 \begin{pmatrix} c \\ 0 \end{pmatrix} + \begin{pmatrix} a \\ b \end{pmatrix} \right] \quad (1.26)$$

$$= \frac{1}{k_2 + 1} \begin{pmatrix} k_2 c + a \\ b \end{pmatrix} \quad (1.27)$$

and

$$F = \frac{k_1 B + A}{k_1 + 1} = \frac{1}{k_1 + 1} \left[k_1 \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \begin{pmatrix} a \\ b \end{pmatrix} \right] \quad (1.28)$$

$$= \frac{1}{k_1 + 1} \begin{pmatrix} a \\ b \end{pmatrix} \quad (1.29)$$

Since O divides BE in the ratio $2 : 1$, using (1.7),

$$O = \frac{2E + B}{3} = \frac{2}{3(k_2 + 1)} \begin{pmatrix} k_2c + a \\ b \end{pmatrix} \quad (1.30)$$

Also, O divides CF in the ratio $2 : 1$, hence,

$$O = \frac{2F + C}{3} = \frac{1}{3} \left[\frac{2}{k_1 + 1} \begin{pmatrix} a \\ b \end{pmatrix} + \begin{pmatrix} c \\ 0 \end{pmatrix} \right] \quad (1.31)$$

$$= \frac{1}{3(k_1 + 1)} \begin{pmatrix} 2a + (k_1 + 1)c \\ 2b \end{pmatrix} \quad (1.32)$$

Equating (1.30) and (1.32),

$$\frac{2}{3(k_2 + 1)} \begin{pmatrix} k_2c + a \\ b \end{pmatrix} = \frac{1}{3(k_1 + 1)} \begin{pmatrix} 2a + (k_1 + 1)c \\ 2b \end{pmatrix} \quad (1.33)$$

$$\Rightarrow \frac{2}{(k_2 + 1)} \begin{pmatrix} k_2c + a \\ b \end{pmatrix} = \frac{1}{(k_1 + 1)} \begin{pmatrix} 2a + (k_1 + 1)c \\ 2b \end{pmatrix} \quad (1.34)$$

Equating the respective coordinates,

$$\frac{2(k_2c + a)}{3(k_2 + 1)} = \frac{2a + (k_1 + 1)c}{k_1 + 1} \quad (1.35)$$

$$\frac{2b}{k_2 + 1} = \frac{2b}{k_1 + 1} \quad (1.36)$$

From (1.36), $k_1 = k_2$ is trivially obtained. Substituting this in (1.35),

$$\frac{2(k_1c + a)}{3(k_1 + 1)} = \frac{2a + (k_1 + 1)c}{k_1 + 1} \quad (1.37)$$

$$\Rightarrow 2k_1c + 2a = 2a + (k_1 + 1)c \quad (1.38)$$

$$k_1 = 1 \quad (1.39)$$

after simplification. Thus, $k_1 = k_2 = 1$ and BE and CF are the medians.

Problem 1.7. In Fig. 1.7, let BE and CF be the medians in $\triangle ABC$. Let the line AD pass through O . Show that AD is also a median.

Proof. Substituting $k_2 = 1$ in (1.30),

$$O = \frac{1}{3} \begin{pmatrix} a + c \\ b \end{pmatrix} \quad (1.40)$$

Since D divides BC in the ratio $k_1 : 1$, from (1.7),

$$D = \frac{k_1C + B}{k_1 + 1} = \frac{1}{k_1 + 1} \begin{pmatrix} c \\ 0 \end{pmatrix} \quad (1.41)$$




Fig. 1.7: Medians of a triangle are concurrent.

Similarly,

$$O = \frac{k_2D + A}{k_2 + 1} = \frac{1}{k_2 + 1} \left[\frac{k_2}{k_1 + 1} \begin{pmatrix} c \\ 0 \end{pmatrix} + \begin{pmatrix} a \\ b \end{pmatrix} \right] \quad (1.42)$$

$$= \frac{1}{(k_2 + 1)(k_1 + 1)} \begin{pmatrix} k_2c + (k_1 + 1)a \\ (k_1 + 1)b \end{pmatrix} \quad (1.43)$$

Equating (1.40) and (1.43),

$$\frac{1}{3} \begin{pmatrix} a + c \\ b \end{pmatrix} = \frac{1}{(k_2 + 1)(k_1 + 1)} \begin{pmatrix} k_2c + (k_1 + 1)a \\ (k_1 + 1)b \end{pmatrix} \quad (1.44)$$

resulting in

$$\frac{a + c}{3} = \frac{k_2c + (k_1 + 1)a}{(k_2 + 1)(k_1 + 1)} \quad (1.45)$$

$$\frac{b}{3} = \frac{(k_1 + 1)b}{(k_2 + 1)(k_1 + 1)} \quad (1.46)$$

after equating the respective coordinates. From (1.46) $k_2 = 2$ is trivially obtained. Substituting $k_2 = 2$ in (1.45),

$$\frac{a + c}{3} = \frac{2c + (k_1 + 1)a}{3(k_1 + 1)} \quad (1.47)$$

$$\Rightarrow (a + c)(k_1 + 1) = 2c + (k_1 + 1)a \quad (1.48)$$

$$\Rightarrow k_1 = 1 \quad (1.49)$$

after simplification. Thus D is the midpoint of BC and AD is also a median. All medians intersect at the point O . This point is known as the centroid.

2 ALTITUDES OF A TRIANGLE

2.1 Lines and Angles

Problem 2.1. In Fig. 2.1, the line passing through the points A, B, C makes an angle θ with the X -axis.

Show that the equation of the line is

$$y = mx + c, \quad (2.1)$$

where (x, y) is any point on the line and $m = \tan \theta$. m is also known as the slope of the line.




Fig. 2.1: The straight line.

Proof. Let $P = (x, y)$ be any point on the line. From the figure, it is obvious that

$$\frac{PR}{CR} = \frac{y - c}{x} = \tan \theta = m \quad (2.2)$$

Rearranging terms,

$$y = mx + c \quad (2.3)$$

Problem 2.2. Show that the equation of the line in Fig. 2.1 can also be expressed as

$$\frac{y - b_2}{x - b_1} = \frac{a_2 - b_2}{a_1 - b_1} \quad (2.4)$$

Proof. Looking at the line segments AB and PB and following the approach in Problem 2.1,

$$\tan \theta = \frac{a_2 - b_2}{a_1 - b_1} = \frac{y - b_2}{x - b_1} \quad (2.5)$$

Problem 2.3. Show that the equation

$$px + qy + r = 0 \quad (2.6)$$

is that of a straight line.

Proof. The above equation can be rearranged as

$$y = -\frac{p}{q}x - \frac{r}{q} \quad (2.7)$$

which is the same as (2.1) with $m = -\frac{p}{q}$ and $c = -\frac{r}{q}$.

Problem 2.4. The point of intersection of the lines

$$p_1x + q_1y + r_1 = 0 \quad (2.8)$$

$$p_2x + q_2y + r_2 = 0 \quad (2.9)$$

is

$$x = -\frac{\begin{vmatrix} r_1 & q_1 \\ r_2 & q_2 \end{vmatrix}}{\begin{vmatrix} p_1 & q_1 \\ p_2 & q_2 \end{vmatrix}} \quad (2.10)$$

$$y = -\frac{\begin{vmatrix} p_1 & r_1 \\ p_2 & r_2 \end{vmatrix}}{\begin{vmatrix} p_1 & q_1 \\ p_2 & q_2 \end{vmatrix}} \quad (2.11)$$

Problem 2.5. Show that angle between the lines

$$y = m_1x + c_1 \quad (2.12)$$

$$y = m_2x + c_2 \quad (2.13)$$

is

$$\theta = \tan^{-1} \frac{m_1 - m_2}{1 + m_1m_2} \quad (2.14)$$

Proof. If θ_1 and θ_2 are the angles made by the two lines with the X-axis respectively, the angle between the two is $\theta = \theta_1 - \theta_2$. Thus, using the trigonometric identity,

$$\tan \theta = \tan(\theta_1 - \theta_2) = \frac{\tan \theta_1 - \tan \theta_2}{1 + \tan \theta_1 \tan \theta_2} \quad (2.15)$$



$$= \frac{m_1 - m_2}{1 + m_1m_2} \quad (2.16)$$

Problem 2.6. Show that the angle between two lines is 90° if $m_1m_2 = -1$.

2.2 Altitude

Definition 2.7. In $\triangle ABC$ Fig. 2.7, $AD \perp BC$ is known as the altitude.

Problem 2.8. In Fig. 2.8, the altitudes AD and BE meet at O . CF is the line obtained by extending CO to meet AB at F . Show that CF is also an altitude of $\triangle ABC$.

Fig. 2.7: Altitude AD .Fig. 2.8: Altitudes AD and BE meet at O .

Proof. The equation for AD is

$$x = a \quad (2.17)$$

The slope of AC is

$$\frac{b}{a - c}. \quad (2.18)$$

Thus, using Problem 2.6, the slope of CE is

$$m_{CE} = \frac{c - a}{b} \quad (2.19)$$

Using (2.1) and (2.19), the equation of CE is

$$y = m_{CE}(x - 0) = \frac{c - a}{b}x \quad (2.20)$$

since O is the intersection of AD and CE , from (2.20) and (2.17), its coordinates are

$$\left(a, \left(\frac{c - a}{b} \right) a \right) \quad (2.21)$$

From (2.21), the slope of CO (and CF) is

$$m_{CF} = \frac{\left(\frac{c - a}{b} \right) a}{a - c} = -\frac{a}{b} \quad (2.22)$$

From Fig. 2.8, the slope of AB is

$$m_{AB} = \frac{b}{a} \quad (2.23)$$

Since


$$m_{AB}m_{CF} = -1, \quad (2.24)$$

using Problem 2.6, $AB \perp CF$ and CF is an altitude of $\triangle ABC$.

3 THE CIRCLE

3.1 Definitions

Definition 3.1. (Circle): Fig. 3.1 shows a circle with centre O and radius r , where each point on the circle is at the same distance r from the centre O .

Fig. 3.1: Circle with radius r .

Problem 3.2. Show that the equation of the circle is

$$(x - a)^2 + (y - b)^2 = r^2 \quad (3.1)$$

Proof. Since each point (x, y) on the circle is at a distance r from the centre (a, b) , using (1.1), (3.1) is obtained.

Definition 3.3. (Tangent): In Fig. 3.3, the line BX that touches the circle at exactly one point C is known as the tangent.

3.2 Tangent

Problem 3.4. In Fig. 3.3, show that the tangent $BX \perp OC$.





Fig. 3.3: Tangent to the circle.

Fig. 3.5: $BP \cdot BQ = BC^2$.

Proof. Since C lies on the circle and is at distance r from the centre O , using (1.1),

$$(x - a)^2 + b^2 = r^2 \quad (3.2)$$

$$\Rightarrow x = a \pm \sqrt{r^2 - b^2} \quad (3.3)$$

By the definition of a tangent, the point C should be unique. Thus, in (3.3), $x = a$ and $r = b$. This implies that $\triangle OCB$ is right angled and $BX \perp OC$.

Problem 3.5. In Fig. 3.5, BX is a tangent to the circle centred at O and touches it at C . P and Q are two points on the circle. Show that

$$BP \cdot BQ = BC^2 \quad (3.4)$$

Proof. Let m be the slope of the line BQ . Then the coordinates of P and Q can be represented as (p, mp) and (q, mq) respectively. The radius of the circle is b . Using (3.1) for P and Q ,

$$(p - a)^2 + (mp - b)^2 = b^2 \quad (3.5)$$

$$\Rightarrow a^2 - 2ap + m^2 p^2 = 2mpb \quad (3.6)$$

and

$$\Rightarrow a^2 - 2aq + m^2 q^2 = 2mqb \quad (3.7)$$

Dividing (3.6) by (3.7),

$$\frac{a^2 - 2ap + p^2(1 + m^2)}{a^2 - 2aq + q^2(1 + m^2)} = \frac{p}{q} \quad (3.8)$$

which, after cross multiplying, results in

$$\begin{aligned} q(a^2 - 2ap + p^2(1 + m^2)) \\ = p(a^2 - 2aq + q^2(1 + m^2)) \end{aligned} \quad (3.9)$$

and can be simplified as

$$a^2(p - q) = pq(p - q)(1 + m^2) \quad (3.10)$$

$$\Rightarrow a^2 = pq(1 + m^2) \quad (3.11)$$

$$\Rightarrow a^2 = p\sqrt{(1 + m^2)} \cdot q\sqrt{(1 + m^2)} \quad (3.12)$$

$$\Rightarrow BC^2 = BP \cdot BQ, \quad (3.13)$$

using (1.1) for obtaining the length of BP and BQ .