

G V V Sharma*

CONTENTS

1	The Right Angled Triangle	1
1.1	Sum of Angles	1
1.2	Budhayana Theorem	2
2	Medians of a Triangle	2
2.1	Area of a Triangle	2
2.2	Median	3
2.3	Similar Triangles	5
3	Angle and Perpendicular Bisectors	7
3.1	Angle Bisectors	7
3.2	Congruent Triangles	7
3.3	Perpendiculars from Vertex to Opposite Side	8
4	Circle	8
4.1	Chord of a Circle	8
4.2	Chords of a circle	8
5	Area of a Circle	10
5.1	The Regular Polygon	10




Fig. 1: Right Angled Triangle




Fig. 2: Sum of angles of a triangle

1 THE RIGHT ANGLED TRIANGLE

Definition 1.1. A right angled triangle looks like Fig. 1. with angles $\angle A$, $\angle B$ and $\angle C$ and sides a , b and c . The unique feature of this triangle is $\angle C$ which is defined to be 90° .

Definition 1.2. For simplicity, let the greek letter $\theta = \angle B$. We have the following definitions.

$$\begin{aligned}
 \sin \theta &= \frac{a}{c} & \cos \theta &= \frac{b}{c} \\
 \tan \theta &= \frac{a}{b} & \cot \theta &= \frac{b}{a} \\
 \csc \theta &= \frac{c}{a} & \sec \theta &= \frac{c}{b}
 \end{aligned}
 \tag{1.1}$$

1.1 Sum of Angles

Definition 1.3. In Fig. 2, the sum of all the angles on the top or bottom side of the straight line XY is 180° .

Definition 1.4. In Fig. 2, the straight line making an angle of 90° to the side AC is said to be parallel to the side BC . Note there is an angle at A that is equal to θ . This is one property of parallel lines. Thus, $\angle YAZ = 90^\circ$.

*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadevall@iith.ac.in.

Problem 1.1. Show that $\angle VAZ = 90^\circ - \theta$

Proof. Considering the line XAZ,

$$\theta + 90^\circ + \angle VAZ = 180^\circ \quad (1.2)$$

$$\Rightarrow \angle VAZ = 90^\circ - \theta \quad (1.3)$$

Problem 1.2. Show that $\angle BAC = 90^\circ - \theta$.

Proof. Consider the line VAB and use the approach in the previous problem. Note that this implies that $\angle VAZ = \angle BAC$. Such angles are known as vertically opposite angles.

Problem 1.3. Sum of the angles of a triangle is equal to 180°

1.2 Budhayana Theorem




Fig. 3: Budhayana Theorem

Problem 1.4. Using Fig. 1, show that

$$\cos \theta = \sin(90^\circ - \theta) \quad (1.4)$$

Proof. From Problem 1.2 and (1.1)

$$\cos(90^\circ - \theta) = \frac{b}{c} = \sin \theta \quad (1.5)$$

Problem 1.5. Using Fig. 3, show that

$$c = a \cos \theta + b \sin \theta \quad (1.6)$$

Proof. We observe that

$$BD = a \cos \theta \quad (1.7)$$

$$AD = b \cos(90 - \theta) = b \sin \theta \quad (\text{From } (1.2)) \quad (1.8)$$

Thus,

$$BD + AD = c = a \cos \theta + b \sin \theta \quad (1.9)$$

Problem 1.6. From (1.6), show that

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (1.10)$$

Proof. Dividing both sides of (1.6) by c ,

$$1 = \frac{a}{c} \cos \theta + \frac{b}{c} \sin \theta \quad (1.11)$$

$$\Rightarrow \sin^2 \theta + \cos^2 \theta = 1 \quad (\text{from } (1.1)) \quad (1.12)$$

Problem 1.7. Using (1.6), show that

$$c^2 = a^2 + b^2 \quad (1.13)$$

(1.13) is known as the Budhayana theorem. It is also known as the Pythagoras theorem.

Proof. From (1.6),

$$c = a \frac{a}{c} + b \frac{b}{c} \quad (\text{from } (1.1)) \quad (1.14)$$

$$\Rightarrow c^2 = a^2 + b^2 \quad (1.15)$$

2 MEDIANS OF A TRIANGLE

2.1 Area of a Triangle




Fig. 4: Area of a Right Triangle

Definition 2.1. The area of the rectangle ACBD shown in Fig. 2.1 is defined as ab . Note that all the angles in the rectangles are 90°

Definition 2.2. The area of the two triangles constituting the rectangle is the same.

Definition 2.3. The area of the rectangle is the sum of the areas of the two triangles inside.

Problem 2.1. Show that the area of ΔABC is $\frac{ab}{2}$

Proof. From (2.3),

$$ar(ABCD) = ar(ACB) + ar(ADB) \quad (2.1)$$

Also from (2.2),

$$ar(ACB) = ar(ADB) \quad (2.2)$$

From (2.1) and (2.2),

$$2ar(ACB) = ar(ABCD) = ab \text{ (from (2.1))} \quad (2.3)$$

$$\Rightarrow ar(ACB) = \frac{ab}{2} \quad (2.4)$$




Fig. 5: Area of a Triangle

Problem 2.2. Show that the area of ΔABC in Fig. 5 is $\frac{1}{2}ah$.

Proof. In Fig. 5,

$$ar(\Delta ADC) = \frac{1}{2}hy \quad (2.5)$$

$$ar(\Delta ADB) = \frac{1}{2}hx \quad (2.6)$$

Thus,

$$ar(\Delta ABC) = ar(\Delta ADC) + ar(\Delta ADB) \quad (2.7)$$

$$= \frac{1}{2}hy + \frac{1}{2}hx = \frac{1}{2}h(x+y) \quad (2.8)$$

$$= \frac{1}{2}ah \quad (2.9)$$

Problem 2.3. Show that the area of ΔABC in Fig. 5 is $\frac{1}{2}ab \sin C$.

Proof. We have

$$ar(\Delta ABC) = \frac{1}{2}ah = \frac{1}{2}ab \sin C \quad (\because h = b \sin C). \quad (2.10)$$

Problem 2.4. Show that

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} \quad (2.11)$$

Proof. Fig. 5 can be suitably modified to obtain

$$ar(\Delta ABC) = \frac{1}{2}ab \sin C = \frac{1}{2}bc \sin A = \frac{1}{2}ca \sin B \quad (2.12)$$

Dividing the above by abc , we obtain

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} \quad (2.13)$$

This is known as the sine formula.

2.2 Median




Fig. 6: Median of a Triangle

Definition 2.4. The line AD in Fig. 6 that divides the side a in two equal halves is known as the median.

Problem 2.5. Show that the median AD in Fig. 6 divides ΔABC into triangles ADB and ADC that have equal area.

Proof. We have

$$ar(\Delta ADB) = \frac{1}{2} \cdot \frac{a}{2} \cdot c \sin B = \frac{1}{4}ac \sin B \quad (2.14)$$

$$ar(\Delta ADC) = \frac{1}{2} \cdot \frac{a}{2} \cdot b \sin C = \frac{1}{4}ab \sin C \quad (2.15)$$

Using the sine formula, $b \sin C = c \sin B$,

$$ar(\Delta ADB) = ar(\Delta ADC) \quad (2.16)$$

Problem 2.6. BE and CF are the medians in Fig. 7. Show that

$$ar(\Delta BFC) = ar(\Delta BEC) \quad (2.17)$$

Proof. Since BE and CF are the medians,

$$ar(\Delta BFC) = \frac{1}{2}ar(\Delta ABC) \quad (2.18)$$

$$ar(\Delta BEC) = \frac{1}{2}ar(\Delta ABC) \quad (2.19)$$

From the above, we infer that

$$ar(\Delta BFC) = ar(\Delta BEC) \quad (2.20)$$




Fig. 7: O is the Intersection of Two Medians

area of a triangle in Fig. 8,

$$\frac{1}{2} \frac{a}{2} AD \sin \theta = \frac{1}{2} \frac{a}{2} AD \sin(180^\circ - \theta) \quad (2.27)$$

$$\Rightarrow \sin \theta = \sin(180^\circ - \theta). \quad (2.28)$$

Note that our geometric definition of $\sin \theta$ holds only for $\theta < 90^\circ$. (2.28) allows us to extend this definition for $\angle ADC > 90^\circ$.




Fig. 8: $\sin \theta = \sin(180^\circ - \theta)$

Problem 2.7. The medians BE and CF in Fig. 7 meet at point O . Show that

$$\frac{OB}{OE} = \frac{OC}{OF} \quad (2.21)$$

Proof. From Problem 2.6,

$$ar(\Delta BFC) = ar(\Delta BEC) \quad (2.22)$$

$$\begin{aligned} \Rightarrow ar(\Delta BOF) + ar(\Delta BOC) \\ = ar(\Delta BOC) + ar(\Delta COE) \end{aligned} \quad (2.23)$$

resulting in

$$ar(\Delta BOF) = ar(\Delta COE) \quad (2.24)$$

Using the sine formula for area of a triangle, the above equation can be expressed as

$$\frac{1}{2} OB OF \sin \theta = \frac{1}{2} OC OE \sin \theta \quad (2.25)$$

$$\Rightarrow \frac{OB}{OE} = \frac{OC}{OF} \quad (2.26)$$

Definition 2.5. We know that the median of a triangle divides it into two triangles with equal area. Using this result along with the sine formula for the




Fig. 9: O divides medians in the ratio 2 : 1

Problem 2.8. In Fig. 9, show that

$$ar(\Delta AFE) = \frac{1}{4}ar(\Delta ABC) \quad (2.29)$$

Proof. We have

$$ar(\Delta AFE) = \frac{1}{2} \frac{b}{2} \frac{c}{2} \sin A = \frac{1}{4} \cdot \frac{1}{2} bc \sin A \quad (2.30)$$

$$= \frac{1}{4} ar(\Delta ABC) \quad (2.31)$$

Problem 2.9. Using Fig. 9, show that

$$\frac{OB}{OE} = \frac{OC}{OF} = 2 \quad (2.32)$$

Proof. Using the sine formula and (2.8), areas of some triangles in Fig. 9 are listed in the following table

Triangle	Area
OFE	$\frac{1}{2}pq \sin \theta$
BOF	$\frac{k}{2}pq \sin \theta$
COE	$\frac{k}{2}pq \sin \theta$
BOC	$\frac{k^2}{2}pq \sin \theta$
BOC	$\frac{1}{2}ar(\Delta ABC)$

where we have used the fact that

$$\sin \angle BOC = \sin(180^\circ - \theta) = \sin \theta \quad (2.33)$$

Since BE is the median

$$\begin{aligned} ar(\Delta BEC) &= \frac{1}{2}ar(\Delta ABC) \\ &= ar(\Delta BOC) + ar(\Delta COE) \\ ar(\Delta BEA) &= \frac{1}{2}ar(\Delta ABC) \\ &= ar(\Delta AFE) + ar(\Delta BOF) + ar(\Delta FOE) \end{aligned} \quad (2.34)$$

Substituting the respective values from the above table,

$$\begin{aligned} \frac{1}{2}ar(\Delta ABC) &= \frac{k^2}{2}pq \sin \theta + \frac{k}{2}pq \sin \theta \\ \frac{1}{2}ar(\Delta ABC) &= \frac{k}{2}pq \sin \theta + \frac{1}{2}pq \sin \theta + \frac{1}{4}ar(\Delta ABC) \end{aligned} \quad (2.35)$$

Simplifying the above,

$$k(k+1) = 2(k+1) \quad (2.36)$$

Since $k \neq -1$, $k = 2$ and the proof is complete.

2.3 Similar Triangles




Fig. 10: The cosine formula

Problem 2.10. In Fig. 10, show that

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} \quad (2.37)$$

Proof. From the figure, the first of the following equations

$$a = b \cos C + c \cos B \quad (2.38)$$

$$b = c \cos A + a \cos C \quad (2.39)$$

$$c = b \cos A + a \cos B \quad (2.40)$$

is obvious and the other two can be similarly obtained. The above equations can be expressed in matrix form as

$$\begin{pmatrix} 0 & c & b \\ c & 0 & a \\ b & a & 0 \end{pmatrix} \begin{pmatrix} \cos A \\ \cos B \\ \cos C \end{pmatrix} = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \quad (2.41)$$

Using the properties of determinants,

$$\cos A = \frac{\begin{vmatrix} a & c & b \\ b & 0 & a \\ c & a & 0 \end{vmatrix}}{\begin{vmatrix} 0 & c & b \\ c & 0 & a \\ b & a & 0 \end{vmatrix}} = \frac{ab^2 + ac^2 - a^3}{abc + abc} = \frac{b^2 + c^2 - a^2}{2abc} \quad (2.42)$$




Fig. 11: Similar Triangles

Problem 2.11. In Fig. 11, show that $EF = \frac{a}{2}$.

Proof. Using the cosine formula for $\triangle AEF$,

$$EF^2 = \left(\frac{b}{2}\right)^2 + \left(\frac{c}{2}\right)^2 - 2\left(\frac{b}{2}\right)\left(\frac{c}{2}\right)\cos A \quad (2.43)$$

$$= \frac{b^2 + c^2 - 2bc \cos A}{4} \quad (2.44)$$

$$= \frac{a^2}{4} \quad (2.45)$$

$$\Rightarrow EF = \frac{a}{2} \quad (2.46)$$

Definition 2.6. The ratio of sides of triangles AEF and ABC is the same. Such triangles are known as similar triangles.

Problem 2.12. Show that similar triangles have the same angles.

Proof. Use cosine formula and the proof is trivial.

Problem 2.13. Show that in Fig. 11, $EF \parallel BC$.

Proof. Since $\triangle AEF \sim \triangle ABC$, $\angle AEF = \angle ACB$. Hence the line $EF \parallel BC$




Fig. 12: Similar Triangles

Problem 2.14. In Fig. 12, the line AO cuts EF at G and is extended to meet the side BC at D . Show that

$$\frac{OA}{OD} = 2. \quad (2.47)$$

Proof. Since $EF \parallel BC$,

$$\triangle AGE \sim \triangle ADC \Rightarrow AG = GD = \frac{AD}{2} \quad (2.48)$$

Also,

$$\triangle OGE \sim \triangle ODB \Rightarrow OD = 2r \quad (2.49)$$

Thus, we have the following relations

$$GD = OG + OD = r + 2r = 3r \quad (2.50)$$

$$AG = GD = 3r \quad (2.51)$$

$$OA = OG + AG = r + 3r = 4r \quad (2.52)$$

Hence

$$\frac{OA}{OD} = \frac{4r}{2r} = 2 \quad (2.53)$$




Fig. 13: Medians meet at a point

Problem 2.15. In Fig. 13, BE is a median of $\triangle ABC$ and $\frac{OA}{OD} = 2$. Show that AD is also a median.

Proof. Using the cosine formula for $\triangle ODE$ and $\triangle OAB$,

$$DE^2 = p^2 + t^2 - 2pq \cos \theta \quad (2.54)$$

$$c^2 = (2p)^2 + (2t)^2 - 2(2p)(2q) \cos \theta \quad (2.55)$$

$$\Rightarrow DE = \frac{c}{2}. \quad (2.56)$$

Now using the cosine formula in $\triangle CDE$ and $\triangle ABC$,

$$\left(\frac{c}{2}\right)^2 = \left(\frac{b}{2}\right)^2 + CD^2 - 2CD\left(\frac{b}{2}\right)\cos C \quad (2.57)$$

$$c^2 = b^2 + a^2 - 2ab \cos C, \quad (2.58)$$

which can be simplified to obtain

$$a^2 - 2ab \cos C = (2CD)^2 - 2(2CD)b \cos C \quad (2.59)$$

$$\Rightarrow (2CD)^2 - 2(2CD)b \cos C + (b \cos C)^2 = a^2 - 2ab \cos C + (b \cos C)^2 \quad (2.60)$$

$$\Rightarrow (2CD - b \cos C)^2 = (a - b \cos C)^2 \quad (2.61)$$

$$\Rightarrow 2CD - b \cos C = a - 2b \cos C \quad (2.62)$$

$$\Rightarrow CD = \frac{a}{2} \quad (2.63)$$

Thus, AD is a median of $\triangle ABC$. *Conclusion:* The medians of a triangle meet at a point.

3 ANGLE AND PERPENDICULAR BISECTORS

3.1 Angle Bisectors




Fig. 14: Angle bisectors meet at a point

Definition 3.1. In Fig. 14, OB divides the $\angle B$ into half, i.e.

$$\angle OBC = \angle OBA \quad (3.1)$$

OB is known as an angle bisector.

OB and OC are angle bisectors of angles B and C . OA is joined and OD, OF and OE are perpendiculars to sides a, b and c .

Problem 3.1. Show that $OD = OE = OF$.

Proof. In $\triangle ODC$ and OEC ,

$$OD = OC \sin \frac{C}{2} \quad (3.2)$$

$$OE = OC \sin \frac{C}{2} \quad (3.3)$$

$$\Rightarrow OD = OE. \quad (3.4)$$

Similarly,

$$OD = OF. \quad (3.5)$$

Problem 3.2. Show that OA is the angle bisector of $\angle A$

Proof. In $\triangle OFA$ and OEA ,

$$OF = OE \quad (3.6)$$

$$\Rightarrow OA \sin OAF = OA \sin OAE \quad (3.7)$$

$$\Rightarrow \sin OAF = \sin OAE \quad (3.8)$$

$$\Rightarrow \angle OAF = \angle OAE \quad (3.9)$$

which proves that OA bisects $\angle A$. *Conclusion:* The angle bisectors of a triangle meet at a point.

3.2 Congruent Triangles

Problem 3.3. Show that in $\triangle ODC$ and OEC , corresponding sides and angles are equal.

Definition 3.2. Note that $\triangle ODC$ and OEC are known as congruent triangles. To show that two triangles are congruent, it is sufficient to show that some angles and sides are equal.

Problem 3.4. SSS: Show that if the corresponding sides of three triangles are equal, the triangles are congruent.

Problem 3.5. ASA: Show that if two angles and any one side are equal in corresponding triangles, the triangles are congruent.

Problem 3.6. SAS: Show that if two sides and the angle between them are equal in corresponding triangles, the triangles are congruent.

Problem 3.7. RHS: For two right angled triangles, if the hypotenuse and one of the sides are equal, show that the triangles are congruent.




Fig. 15: Perpendicular bisectors meet at a point

Definition 3.3. In Fig. 15, $OD \perp BC$ and $BD = DC$. OD is defined as the perpendicular bisector of BC .

Problem 3.8. In Fig. 15, show that $OA = OB = OC$.

Proof. In $\triangle ODB$ and ODC , using Budhayana's theorem,

$$\begin{aligned} OB^2 &= OD^2 + BD^2 \\ OC^2 &= OD^2 + DC^2 \end{aligned} \quad (3.10)$$

Since $BD = DC = \frac{a}{2}$, $OB = OC$. Similarly, it can be shown that $OA = OC$. Thus, $OA = OB = OC$.

Definition 3.4. In $\triangle AOB$, $OA = OB$. Such a triangle is known as an isosceles triangle.

Problem 3.9. Show that $AF = BF$.

Proof. Trivial using Budhayana's theorem. This shows that OF is a perpendicular bisector of AB .
Conclusion: The perpendicular bisectors of a triangle meet at a point.

3.3 Perpendiculars from Vertex to Opposite Side




Fig. 16: Perpendiculars from vertex to opposite side meet at a point

In Fig. 16, $AD \perp BC$ and $BE \perp AC$. CF passes through O and meets AB at F .

Problem 3.10. Show that

$$OE = c \cos A \cot C \quad (3.11)$$

Proof. In Δ s AEB and AEO ,

$$AE = c \cos A \quad (3.12)$$

$$OE = AE \tan(90^\circ - C) (\because ADC \text{ is right angled}) \quad (3.13)$$

$$= AE \cot C \quad (3.14)$$

From both the above, we get the desired result.

Problem 3.11. Show that $\alpha = A$.

Proof. In ΔOEC ,

$$CE = a \cos C (\because BEC \text{ is right angled}) \quad (3.15)$$

Hence,

$$\begin{aligned} \tan \alpha &= \frac{CE}{OE} \\ &= \frac{a \cos C}{c \cos A \cot C} \\ &= \frac{a \cos C \sin C}{c \cos A \cos C} \\ &= \frac{a \sin C}{c \cos A} \\ &= \frac{c \sin A}{c \cos A} \left(\because \frac{a}{\sin A} = \frac{c}{\sin C} \right) \\ &= \tan A \end{aligned} \quad (3.16)$$

$$\Rightarrow \alpha = A$$

Problem 3.12. Show that $CF \perp AB$

Proof. Consider triangle OFB and the result of the previous problem. \therefore the sum of the angles of a triangle is 180° , $\angle CFB = 90^\circ$.
Conclusion: The perpendiculars from the vertex of a triangle to the opposite side meet at a point.

4 CIRCLE

4.1 Chord of a Circle




Fig. 17: Circle Definitions

Definition 4.1. Fig. 17 represents a circle. The points in the circle are at a distance r from the centre O . r is known as the radius.

4.2 Chords of a circle

Definition 4.2. In Fig. 17, A and B are points on the circle. The line AB is known as a chord of the circle.

Problem 4.1. In Fig. 18 Show that $\angle OAB = 2\angle APB$.




Fig. 18: Angle subtended by chord AB at the centre O is twice the angle subtended at P .

Proof. In Fig. 18, the triangles OPA and OPB are isosceles. Hence,

$$\angle OPB = \angle OBP = \theta_1 \tag{4.1}$$

$$\angle OPA = \angle OAP = \theta_2 \tag{4.2}$$

Also, α and β are exterior angles corresponding to the triangle OPB and OPA respectively. Hence

$$\alpha = 2\theta_1 \tag{4.3}$$

$$\beta = 2\theta_2 \tag{4.4}$$

Thus,

$$\angle AOB = \alpha + \beta \tag{4.5}$$

$$= \theta_1 + \theta_2 \tag{4.6}$$

$$= \angle APB \tag{4.7}$$

Definition 4.3. The diameter of a circle is the chord that divides the circle into two equal parts. In Fig. 19, AB is the diameter and passes through the centre O




Fig. 19: Diameter of a circle.

Problem 4.2. In Fig. 19, show that $\angle APB = 90^\circ$.




Fig. 20: $PA.PB = PC.PD$

Problem 4.3. In Fig. 20, show that

$$\begin{aligned} \angle ABD &= \angle ACD \\ \angle CAB &= \angle CDB \end{aligned} \tag{4.8}$$

Proof. Use Problem 4.1.

Problem 4.4. In Fig. 20, show that the triangles PAB and PBD are similar

Proof. Trivial using previous problem

Problem 4.5. In Fig. 20, show that

$$PA.PB = PC.PD \tag{4.9}$$

Proof. Since triangles PAC and PBD are similar,

$$\frac{PA}{PD} = \frac{PC}{PB} \tag{4.10}$$

$$\Rightarrow PA.PB = PC.PD \tag{4.11}$$




Fig. 21: Tangent to a Circle.

Definition 4.4. The line PX in Fig. 21 touches the circle at exactly one point P . It is known as the tangent to the circle.




Fig. 22: Shortest distance from O to line PX

Problem 4.6. OP is the perpendicular to the line PX as shown in the Fig. 22. Show that OP is the shortest distance between the point O and the line PX .

Proof. Let P_1 be a point on the line PX . Then OPP_1 is a right angled triangle. Using Budhayana's theorem,

$$OP_1^2 = OP^2 + PP_1^2 \quad (4.12)$$

$$\Rightarrow OP_1 > OP$$

Thus, OP is the shortest distance between O and line PX .

Problem 4.7. Show that $\angle OPX = 90^\circ$

Proof. In Fig. 21, we can see that OP is the radius of the circle and the length of all line segments from O to the line $PX > r$. Using the result of the previous problem, it is obvious that $OP \perp PX$.




Fig. 23: $PA.PB = PC^2$.

Problem 4.8. In Fig. 23 show that

$$\angle PCA = \angle PBC \quad (4.13)$$

O is the centre of the circle and PC is the tangent.

Proof. For convenience, greek letters are used for representing certain angles. Since ΔOAC is isosceles,

$$2\alpha + 2(\beta - \phi) = 180^\circ \quad (4.14)$$

$$\Rightarrow \alpha + (\beta - \phi) = 90^\circ \quad (4.15)$$

$$\Rightarrow \alpha + \beta = 90^\circ + \phi \quad (4.16)$$

Since θ is an exterior angle for the ΔABC ,

$$\theta = \alpha + \beta \quad (4.17)$$

From both the above equations

$$\theta = 90^\circ + \phi \quad (4.18)$$

Since PC is the tangent,

$$\angle PCB = 90^\circ + \phi = \theta \quad (4.19)$$

Considering the sum of angles in ΔPAC ΔPBC ,

$$\angle P + \theta + \angle PCA = 180^\circ \quad (4.20)$$

$$\angle P + \theta + \alpha = 180^\circ \quad (4.21)$$

Hence,

$$\angle PCA = \alpha \quad (4.22)$$

Problem 4.9. In Fig. 23, show that the triangles PAC and PBC are similar.

Proof. From the previous problem, it is obvious that corresponding angles of both triangles are equal. Hence they are similar.

Problem 4.10. Show that $PA.PB = PC^2$

Proof. Since $\Delta PAC \sim \Delta PBC$, their sides are in the same ratio. Hence,

$$\frac{PA}{PC} = \frac{PC}{PB} \quad (4.23)$$

$$\Rightarrow PA.PB = PC^2 \quad (4.24)$$

Problem 4.11. In Fig. 24, show that

$$PA.PB = PC.PD \quad (4.25)$$

Proof. Draw a tangent and use the previous problem.

5 AREA OF A CIRCLE

5.1 The Regular Polygon

Definition 5.1. In Fig. 25, 6 congruent triangles are arranged in a circular fashion. Such a figure is




Fig. 24: $PA \cdot PB = PC^2$.

known as a regular hexagon. In general, n number of triangles can be arranged to form a regular polygon.




Fig. 25: Polygon Definition

Definition 5.2. The angle formed by each of the congruent triangles at the centre of a regular polygon of n sides is $\frac{360^\circ}{n}$.




Fig. 26: Polygon Area

Problem 5.1. Show that the area of a regular polygon is given by

$$\frac{n}{2}r^2 \sin \frac{360^\circ}{n} \tag{5.1}$$

Proof. The triangle that forms the polygon of n sides is given in Fig. 26. Thus,

$$\begin{aligned} ar(\text{polygon}) &= nar(\Delta ABC) \\ &= \frac{n}{2}r^2 \sin \frac{360^\circ}{n} \end{aligned} \tag{5.2}$$




Fig. 27: Circle Area in between Area of Two Polygons

Problem 5.2. Using Fig. 27, show that

$$\frac{n}{2}r^2 \sin \frac{360^\circ}{n} < \text{area of circle} < nr^2 \tan \frac{180^\circ}{n} \tag{5.3}$$

The portion of the circle visible in Fig. 27 is defined to be a sector of the circle.

Proof. Note that the circle is squeezed between the inner and outer regular polygons. As we can see from Fig. 27, the area of the circle should be in between the areas of the inner and outer polygons. Since

$$ar(\Delta OAB) = \frac{1}{2}r^2 \sin \frac{360^\circ}{n} \tag{5.4}$$

$$ar(\Delta OPQ) = 2 \times \frac{1}{2} \times r \tan \frac{360/n}{2} \times r \tag{5.5}$$

$$= r^2 \tan \frac{180^\circ}{n}, \tag{5.6}$$

we obtain (5.3).

Problem 5.3. Using Fig. 28, show that

$$\sin \theta_1 = \sin (\theta_1 + \theta_2) \cos \theta_2 - \cos (\theta_1 + \theta_2) \sin \theta_2 \tag{5.7}$$




Fig. 28: $\sin 2\theta = 2 \sin \theta \cos \theta$

Proof. The following equations can be obtained from the figure using the formula for the area of a triangle

$$ar(\Delta ABC) = \frac{1}{2}ac \sin(\theta_1 + \theta_2) \quad (5.8)$$

$$= ar(\Delta BDC) + ar(\Delta ADB) \quad (5.9)$$

$$= \frac{1}{2}cl \sin \theta_1 + \frac{1}{2}al \sin \theta_2 \quad (5.10)$$

$$= \frac{1}{2}ac \sin \theta_1 \sec \theta_2 + \frac{1}{2}a^2 \tan \theta_2$$

$$(\because l = a \sec \theta_2) \quad (5.11)$$

$$\Rightarrow \sin(\theta_1 + \theta_2) = \sin \theta_1 \sec \theta_2 + \frac{a}{c} \tan \theta_2 \quad (5.12)$$

$$\Rightarrow \sin(\theta_1 + \theta_2) = \sin \theta_1 \sec \theta_2 + \cos(\theta_1 + \theta_2) \tan \theta_2 \quad (5.13)$$

$$\Rightarrow \sin(\theta_1 + \theta_2) \cos \theta_2 = \sin \theta_1 + \cos(\theta_1 + \theta_2) \sin \theta_2 \quad (5.14)$$

$$\Rightarrow \sin \theta_1 = \sin(\theta_1 + \theta_2) \cos \theta_2 - \cos(\theta_1 + \theta_2) \sin \theta_2 \quad (5.15)$$

Problem 5.4. Prove the following identities

1) $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta.$

2) $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta.$

Proof. In (5.7), let

$$\begin{aligned} \theta_1 + \theta_2 &= \alpha \\ \theta_2 &= \beta \end{aligned} \quad (5.16)$$

This gives (5.4). In (5.4), replace α by $90^\circ - \alpha$. This

results in

$$\sin(\alpha - \beta) = \sin(90^\circ - \alpha - \beta) \quad (5.17)$$

$$= \sin(90^\circ - \alpha) \cos \beta - \cos(90^\circ - \alpha) \sin \beta \quad (5.18)$$

$$\Rightarrow \cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \quad (5.19)$$

Problem 5.5. Using (5.7) and (2), show that

$$\sin(\theta_1 + \theta_2) = \sin \theta_1 \cos \theta_2 + \cos \theta_1 \sin \theta_2 \quad (5.20)$$

$$\cos(\theta_1 - \theta_2) = \cos \theta_1 \cos \theta_2 + \sin \theta_1 \sin \theta_2 \quad (5.21)$$

Proof. From (5.7),

$$\sin(\theta_1 + \theta_2) \cos \theta_2 = \sin \theta_1 + \cos(\theta_1 + \theta_2) \sin \theta_2 \quad (5.22)$$

Using (2) in the above,

$$\begin{aligned} \sin(\theta_1 + \theta_2) \cos \theta_2 &= \sin \theta_1 + (\cos \theta_1 \cos \theta_2 \\ &\quad - \sin \theta_1 \sin \theta_2) \sin \theta_2 \end{aligned} \quad (5.23)$$

which can be expressed as

$$\begin{aligned} \sin(\theta_1 + \theta_2) \cos \theta_2 &= \sin \theta_1 + \cos \theta_1 \cos \theta_2 \sin \theta_2 \\ &\quad - \sin \theta_1 \sin^2 \theta_2 \end{aligned} \quad (5.24)$$

Since

$$\sin^2 \theta_2 = 1 - \cos^2 \theta_2, \quad (5.25)$$

we obtain

$$\begin{aligned} \sin(\theta_1 + \theta_2) \cos \theta_2 &= \cos \theta_1 \cos \theta_2 \sin \theta_2 \\ &\quad + \sin \theta_1 \cos^2 \theta_2 \end{aligned} \quad (5.26)$$

resulting in

$$\sin(\theta_1 + \theta_2) = \cos \theta_1 \sin \theta_2 + \sin \theta_1 \cos \theta_2 \quad (5.27)$$

after factoring out $\cos \theta_2$. Using a similar approach, (5.21) can also be proved.

Problem 5.6. Show that

$$\sin 2\theta = 2 \sin \theta \cos \theta \quad (5.28)$$

Problem 5.7. Show that

$$\cos^2 \frac{180^\circ}{n} < \frac{\text{area of circle}}{nr^2 \tan \frac{180^\circ}{n}} < 1 \quad (5.29)$$

Proof. From (5.3) and the previous result,

$$\frac{n}{2} r^2 \sin \frac{360^\circ}{n} < \text{area of circle} < nr^2 \tan \frac{180^\circ}{n} \quad (5.30)$$

Problem 5.16. Show that

$$\Rightarrow nr^2 \sin \frac{180^\circ}{n} \cos \frac{180^\circ}{n} < \text{area of circle} < nr^2 \tan \frac{180^\circ}{n} \quad (5.31)$$

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1 \quad (5.41)$$

Problem 5.8. Show that if

$$\theta_1 < \theta_2, \sin \theta_1 < \sin \theta_2. \quad (5.32)$$

Proof. Trivial using the definition and choosing angles θ_1 and θ_2 appropriately in a right angled triangle.

Problem 5.9. Show that if

$$\theta_1 < \theta_2, \cos \theta_1 > \cos \theta_2. \quad (5.33)$$

Problem 5.10. Show that

$$\sin 0^\circ = 0 \quad (5.34)$$

Proof. Follows from (5.33).

Problem 5.11. Show that

$$\cos 0^\circ = 1 \quad (5.35)$$

Problem 5.12. Show that for large values of n

$$\cos^2 \frac{180^\circ}{n} = 1 \quad (5.36)$$

Proof. Follows from previous problem.

Definition 5.3. The previous result can be expressed as

$$\lim_{n \rightarrow \infty} \cos^2 \frac{180^\circ}{n} = 1 \quad (5.37)$$

Problem 5.13. Show that

$$\text{area of circle} = r^2 \lim_{n \rightarrow \infty} n \tan \frac{180^\circ}{n} \quad (5.38)$$

Definition 5.4.

$$\pi = \lim_{n \rightarrow \infty} n \tan \frac{180^\circ}{n} \quad (5.39)$$

Thus, the area of a circle is πr^2 .

Definition 5.5. The radian is a unit of angle defined by

$$1 \text{ radian} = \frac{360^\circ}{2\pi} \quad (5.40)$$

Problem 5.14. Show that the circumference of a circle is $2\pi r$.

Problem 5.15. Show that the area of a sector with angle θ in radians is $\frac{1}{2}r^2\theta$.